Asociación Española
Enfermedad de Andrade
(Amiloidosis Hereditaria por Transtirretina)

II JORNADAS AMILOIDOSIS HEREDITARIA por TRANSTIRRETINA (AhTTR)

Viernes 5 de Octubre de 2018
Hospital Son Llàtzer (Salón de Actos)
Palma de Mallorca

DIAGNÓSTICO PRECOZ EN LA AH-TTR: “PERSIGUIENDO LA NEUROPATIA DE FIBRA FINA”.

● PRINCIPALES EXPLORACIONES NEUROFISIOLÓGICAS.

Dra. CRISTINA DESCALS MOLL .
NEUROFISIOLOGIA CLINICA.SERVICIO DE NEUROLOGIA.
EQUIPO MULTIDISCIPLINAR AMILOIDOSIS POR TTR.
H.SON LLÀTZER.PALMA DE MALLORCA.

INTRODUCCION:

- Ah-TTR/TTR-FAP:

- Enfermedad sistémica, progresiva, neurodegenerativa, rara, severa, crónica (trat^o), discapacitante, de HAD e inicio en edad adulta.

- Descrita por 1^a vez en 1952 (Dr. Corino de Andrade).

- Focos endémicos

- Portugal, Suecia, Japón, España (Mallorca/Huelva).

- PREVALENCIA GLOBAL: **10.000 personas...** Y SUS FAMILIAS.

- Prevalencia en Portugal : **22.93/100.000h.** (1865 pacientes). (2016).
- Mallorca :95 pacientes (2016).

RESEARCH REPORT

Transthyretin familial amyloid polyneuropathy (TTR-FAP) in Mallorca: a comparison between late- and early-onset disease

Juan Buaides-Reinés¹, Manuel Raya-Cruz¹, Cristina Gallego-Lezain¹, Tomás Ripoll-Vera², Mercedes Udín-Martín³, Hernán Andreu-Serra⁴, and Eugenia Cineseros-Barroso⁵

¹Department of Internal Medicine, Son Llàtzer Hospital, Caixa de Manacor; ²Department of Cardiology, Son Llàtzer Hospital, Caixa de Manacor; ³Department of Neurology, Son Llàtzer Hospital, Caixa de Manacor; ⁴Department of Digestive Medicine, Son Llàtzer Hospital, Caixa de Manacor; and ⁵Clinical Research Coordinator of TTR Group, Son Llàtzer Hospital, Caixa de Manacor, Palma, Spain

Neuro
epidemiology

Original Paper

Neuroepidemiology 2018;51:177–182
DOI: 10.1159/000488023

Received February 14, 2018
Revised April 10, 2018
Published online August 16, 2018

Epidemiology of Transthyretin Familial Amyloid Polyneuropathy in Portugal: A Nationwide Study

Mónica Inês^a, Teresa Coelho^{b,c}, Isabel Conceição^{a,d}, Filipa Duarte-Ramos^a, Mamede de Carvalho^{a,d}, João Costa^{a,e}

^aInstituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal; ^bAndrade's Center for Familial Amyloidosis, Porto, Portugal; ^cDepartment of Neurosciences, Hospital de Santo António, Porto, Portugal; ^dDepartment of Neurosciences and Mental Health, Hospital de Santa Maria, Lisboa, Portugal; ^eDepartment of Social Pharmacy, Faculty of Pharmacy, University of Lisbon, Lisbon, Portugal; ^fLaboratório de Farmacologia Clínica e Terapêutica, Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal

II JORNADAS AMILOIDOSIS HEREDITARIA por TRANSTIRRETINA (A β TTR)

ESTIMATING THE GLOBAL PREVALENCE OF TRANSTHYRETIN FAMILIAL AMYLOID POLYNEUROPATHY

HARTMUT H. SCHMIDT, MD,¹ MARCIA WADDINGTON-CRUZ, MD, PhD,² MARC F. BOTTEMAN, MSc, MA,³ JOHN A. CARTER, MS,⁴ AVIJEET S. CHOPRA, PhD,³ MARKAY HOPPS, MPH,⁵ MICHELLE STEWART, PhD,⁶ SHARI FALLET, DO,⁶ and LESLIE AMASS, PhD⁷

Muscle Nerve 57: 829–837, 2018

FIGURE 2. Extracted ATTR-FAP prevalence values per 1 million general population (1M). The highest and lowest prevalence values reported in the literature were for subnational regions. The highest prevalence was reported in Northern Portugal (1,631.20/1M), and the lowest was reported in Sicily (0.20/1M). Values are categorized as A (endemic country, national estimate), B (endemic country, regional estimate), C (nonendemic country, national estimate), and D (nonendemic country, regional estimate). ATTR-FAP, transthyretin familial amyloid polyneuropathy

Table 2. Prevalence estimates by country

Country	General population, M	Prevalence low	Prevalence mid	Prevalence high
Totals	4,582.3	5,526	10,186	38,468
Core	460.1	3,639	3,762	3884
Turkey	78.7	25	25	25
Bulgaria	7.2	41	41	41
Netherlands	16.9	45	45	45
Cyprus	1.2	51	51	51
Germany	81.4	121	121	121
Japan	127.0	111	123	135
Sweden	9.8	253	253	253
France	66.8	502	502	502
Italy	60.8	500	550	600
Portugal	10.3	1,990	2,051	2,111
Extrapolated	4,122.2	1,887	6,424	3,4584
Luxembourg	0.6	0	1	4
Slovenia	2.1	1	3	16
Ireland (North)	1.9	1	3	14
New Zealand	4.6	1	7	35
Finland	5.5	2	8	41
Denmark	5.7	2	8	43
Israel	8.4	3	12	63
Switzerland	8.3	3	12	62
Austria	8.6	3	13	65
Hungary	9.8	3	15	74
Czech Republic	10.6	3	16	79
Greece	10.8	3	16	81
Belgium	11.3	4	17	85
Ecuador	16.1	5	24	121
Romania	19.8	6	29	149
Sri Lanka	21.0	7	31	158
Taiwan	23.5	8	35	177
Australia	23.8	8	35	179
Korea (South)	50.6	16	75	381
Malaysia	30.3	10	45	228
Canada	35.9	12	53	270
Poland	12	12	56	286
Argentina	14	14	64	326
Spain	46.4	15	69	349
UK	65.1	21	97	490
Mexico	127.0	41	188	955
Russia	144.1	46	214	1,084
Bangladesh	161.0	52	239	1,211
Brazil*	207.8	623	623	5,078
USA†	321.4	104	476	2,488
India	1,311.1	423	1,943	9,858
China	1,347.7	435	1,997	10,134

PUNTOS CLAVE :

- El diagnóstico está frecuentemente retrasado de 2,8-4,3 años TRAS EL PRIMER SÍNTOMA.
- El pronóstico es FATAL entre los 2 a 15 desde el inicio.
- La PNP progresa en 3 a 5 años.
 - Primero FF autonómicas >FF termoalgésica> FG (S>M).
 - Distal a proximal de EEII a EESS.
- Forma de inicio PRECOZ (<50 a.) : predomina la PNP y el inicio con la neuropatía de fibra fina.
- Forma de inicio TARDÍO (>50 a.) : predomina la afectación cardiaca +PNP (más difusa).
- TRATAMIENTOS EMERGENTES : DIAGNÓSTICO PRECOZ.

Clinical features of transthyretin familial amyloid polyneuropathy

Within most of the regions in which it is endemic (Portugal, Cyprus, and Majorca), TTR-FAP occurs before age 40 years (early onset), with progressive sensory-motor and autonomic neuropathy leading to cachexia and eventually death. On average, 10.8 years of disease onset [7,48,59]. Length-dependent small-fibre sensory-motor polyneuropathy with life-threatening autonomic dysfunction is a distinguishing feature of TTR-FAP in these regions, but frequently, there is also cardiac (conduction disturbance), renal, and ocular involvement [60^{***}].

Sixty years of transthyretin familial amyloid polyneuropathy (TTR-FAP) in Europe: where are we now? A European network approach to defining the epidemiology and management patterns for TTR-FAP

Yesim Parman^a, David Adams^b, Laura Obici^c, Lucía Galán^d,
Velina Guergueltcheva^a, Ole B. Suhr^f, Teresa Coelho^g,
from the European Network for TTR-FAP (ATReuNET)

“HABRA QUE BUSCARLA Y DEMOSTRARLA” ...

NUESTRA EXPERIENCIA.

- 👉 ESTUDIO DE SEGUIMIENTO EN PORTADORES VAL30MET EN LA UNIDAD MULTIDISCIPLINAR DE AMILOIDOSIS.
- 👉 PRINCIPALES PRUEBAS NEUROFISIOLÓGICAS.

“PERSIGUIENDO LA NEUROPATIA DE FIBRA FINA”.

TIPOS DE FIBRAS NERVIOSAS

Axones aferentes primarios

Medscape® www.medscape.com

Motor		Sensory		Autonomic	
Myelinated	Myelinated	Thinly myelinated	Un-myelinated	Thinly myelinated	Un-myelinated
A alpha	A alpha/beta	A delta	C	C	C
Large		Small			
Muscle control	Touch, vibration, position perception	Cold perception, Pain	Warm perception, Pain	Heart rate, blood pressure, sweating, GIT, GUT function	

Source: Endocr Pract © 2007 American Association of Clinical Endocrinologists

NEUROPATIA de FIBRA FINA:

- Descrita en 1992: Stewart et al.

The purpose of this study was to evaluate sweating and cardiovascular autonomic function in patients with distal small fiber neuropathy (DSFN). Sweat testing by the quantitative sudomotor axon reflex test was abnormal in 32 of 40 (80%) patients. The thermoregulatory sweat test was abnormal in 18 of 25 (72%) patients; one or both tests were abnormal in 36 of 40 (90%). Minor heart rate abnormalities were present in 11 of 40 (28%) patients. We conclude that, in patients with DSFN: (a) the sympathetic sudomotor fibers are frequently affected, and that evaluation of sweating is a useful diagnostic test; (b) that the autonomic nerves controlling heart rate are less affected.

Key words: small fiber neuropathy • sweat testing • cardiovascular reflexes
MUSCLE & NERVE 15:661-665 1992

DISTAL SMALL FIBER NEUROPATHY: RESULTS OF TESTS OF SWEATING AND AUTONOMIC CARDIOVASCULAR REFLEXES

JOHN D. STEWART FRCP(C), PHILLIP A. LOW, MD,
and ROBERT D. FEALEY, MD

- Aquella neuropatía periférica en la que se afectan las fibras nerviosas de pequeño calibre : A δ y C.
 - La mayoría de PNP afectan en su evolución los dos tipos de fibras (f+g).
 - Los nervios sensitivos están formados por fibras g+f con diferentes umbrales de estímulo y receptores.
 - Si existe > afectación del SNA (fibras amielínicas C):
 - NEUROPATÍA AUTONÓMICA.
 - Si existe > afectación difusa de FF (A δ y C : termoelgésica+autonómica) :
 - NEUROPATÍA DE FIBRA FINA.

NEUROPATIA de FIBRA FINA:

• FISIOPATOLOGÍA:

- Degeración axonal de esas terminaciones nerviosas.
- Hiperexcitabilidad (canales Na⁺).

• ETIOLOGÍA:

- Idiopática.
- Secundaria.

Table 2. Causes of small fiber neuropathy

Etiology

Metabolic/nutritional

- Diabetes mellitus,^{44,45} impaired fasting glucose,^{46,47} and "insulin neuritis"⁵²
- Metabolic syndrome^{58,59}
- Chronic kidney disease⁶⁰
- Vitamin B₁₂ deficiency¹⁴
- Vitamin B₁ deficiency^{54,55}
- Thyroid dysfunction^{61,125}
- **Hyperlipidemia**^{62,63}

Infectious

- Human immunodeficiency virus / acquired immunodeficiency syndrome^{65,66}
- Hepatitis C⁶⁴
- **Chagas disease**⁶⁸

Amyloidosis⁶⁹⁻⁷¹

- Systemic
- Familial

Autoimmune

- **Sjogren syndrome**⁷²⁻⁷⁴
- Fibromyalgia^{125,126}
- Systemic lupus erythematosus⁷⁵
- Vasculitis⁷⁶
- Sarcoidosis⁸³⁻⁸⁵

Drugs and toxins

- Metronidazole⁸⁶
- Nitrofurantoin⁸⁷
- Linezolid^{89,127}
- Anti-TNF inhibitor⁹¹
- Thallium⁸⁸
- Statins⁹³
- Chemotherapeutic agents, such as bortezomib⁹²
- Vaccination for rabies, varicella, or Lyme⁹⁰
- Alcohol^{56,57}

Paraneoplastic

- Multiple myeloma and monoclonal gammopathies⁹⁴
- Lung cancer⁹⁵

Hereditary

- Fabry disease⁹⁶
- Sodium channelopathies (Na_v1.7, Na_v1.8)^{23,106}
- Hereditary sensory and autonomic neuropathy types II, III, IV^{93,128}

Inflammatory

- Sensory/small fiber **Guillain-Barré syndrome**^{100,101}

Other rare associations

- Parkinson disease¹²⁹
- Pompe disease¹³⁰
- Fragile X-associated tremor ataxia syndrome¹³¹
- Benign fasciculation syndrome¹³²
- Critical illness¹³³
- Celiac disease¹³⁴
- Wilson disease^{135,136}
- Amyotrophic lateral sclerosis^{137,138}
- X-linked adrenoleukodystrophy¹³⁹

Idiopathic⁴²

INVITED REVIEW

SMALL FIBER NEUROPATHY: GETTING BIGGER!

AMANDA C. Y. CHAN, MBBS, FRCP,¹ and BINAR P. WILDER-SMITH, MD²

¹ Division of Neurology, National University Hospital, Level 10 Tower Block, University Medicine Cluster, 1E Kent Ridge Road, 119226, Singapore

² Neurology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore

Accepted 10 February 2016

NFF: SINTOMAS y SIGNOS

DOLOR-DISESTESIAS

- Severo, limitante
 - > espontáneo.
 - Evocado (sábanas en la cama, calcetines).
 - Ejercicio o calor (Canalopatías, Fabry).
- Distribución longitud-dependiente (pies>manos).
- Ocasionalmente distribución focal (inmunomediadas).
- Quemazón, pinchazos, prurito.
- HIPOESTESIA (<frecuente).

ALTERACIÓN SENSIBILIDAD A LA TEMPERATURA

- Hipoestesia al frío, calor.
- Hiperalgnesia al frío, calor.

SÍNTOMAS AUTONÓMICOS (<20%)

- Hipotensión ortostática.
- Disfunción gastrointestinal-urinaria.
- Disfunción sexual.
- Ojo seco.

Tabla 2 Síntomas autonómicos reportados en esta cohorte de 88 pacientes con neuropatía de fibra fina

Síntoma autonómico	No. de pacientes (%)
Síntomas síca (ojo seco y/o boca seca)	17 (19,3)
Hiperhidrosis	16 (18,2)
Alteraciones defecatorias	12 (13,6)
Mareo ortostático	10 (11,4)
Alteraciones miccionales	9 (10,2)
Sofocos	5 (5,7)
Disfunción sexual	5 (5,7)
Gastroparenia	2 (2,3)
Alteración en la acomodación	2 (2,3)
Hipohidrosis	1 (1,1)
Palpitaciones	1 (1,1)

- Cambios tróficos piel: seca, agrietada o brillante.
- Normalidad función motora.
- Normalidad función sensitiva de fibra gruesa: tacto, vibración y propiocepción.
- Disminución sensibilidad punzante y térmica.
- Hiperalgnesia al pinchazo.
- Alodinia al pincel.
- Disminución TA cambio postural con aumento FC.

- SPI
- CALAMBRES

II JORNADAS AMILOIDOSIS HEREDITARIA por TRANSTIRRETINA (A β TTR)

Asociación Española
Enfermedad de Andrade
(Amiloidosis Hereditaria por Transtirretina)

The diagnostic challenge of small fibre neuropathy: clinical presentations, evaluations, and causes

Astrid J. Terkelsen*, Pål Karlsson*, Giuseppe Lauria, Roy Freeman, Nanna B. Finnerup, Trods S. Jensen

Lancet Neurol 2017; 16: 934-44

DIAGNÓSTICO: Exploraciones Complementarias

- CUESTIONARIOS PARA VALORAR NFF.
- PRUEBAS VALORACIÓN FUNCIONAL SNA:
 - CARDIOVASCULAR: HRV, Tilt Test, Hipotension postural (TA).
 - MN: SPECT (I-MIBG).
 - PRUEBAS DE VALORACION FUNCION SUDOMOTORA:
 - QSART.
 - SUDOSCAN (ESC).
 - STIMULATED SKIN WRINKLING
 - TEST de SUDOR TERMORREGULADO, MOLDES DE “SWEAT IMPRINT”,...

- PRUEBAS NEUROFISIOLÓGICAS: **ENG por definición debe ser NORMAL.**

- QST.
- RSC.
- Intervalo R-R.
- POTENCIALES EVOCADOS LASER/CALOR (CHEPs /LEPs).
- Microneurografía.

- PRUEBAS DE A.P.: Biopsia cutánea (DFNIE).
- MICROSCOPIA CONFOCAL (Oftalmología).

DIAGNÓSTICO

- **DIAGNÓSTICO de la NEUROPATÍA DE FIBRA FINA:**

- **CLÍNICO** con el soporte de **UNA TÉCNICA DIAGNÓSTICA ESPECÍFICA** para el estudio de **LA FIBRA FINA > TERMOALGÉSICA.**

- Criterios diagnósticos.

- Las exploraciones complementarias para el estudio específico de las fibras de pequeño calibre son **MUY LABORIOSAS** (**“time-consuming”**), y están disponibles en pocos centros hospitalarios.

- PEL/CHEP/microneurografía.

- Unidades de estudio de DISAUTONOMIA.

- Biopsia cutánea para valorar la DFNIE.

CRITERIOS DIAGNÓSTICOS

AL MENOS 2 (2006):

- **1) SINTOMAS CLÍNICOS DE AFECTACIÓN DE FIBRA FINA:**
 - Alodinia y/o hiperalgesia y/o alteración s. termoalgésica.
 - Distribución de neuropatía periférica.
- **2) QST patológico (para umbral de frío y calor).**
- **3) Disminución de la densidad distal EEl de fibras (biopsia piel) : IENF.**

- **DESCARTAR: PNP** (no déficit motor, no alteración sensibilidad vibratoria/ artrocinética/táctil /RMP, no alteración ENG) .

- **2009:** POSIBLE (1), PROBABLE (1+ ENG n.sural NORMAL), DEFINIDA (1+2+biopsia cutánea).

doi:10.1093/brain/awn093

Brain (2008), 131, 1912–1925

The diagnostic criteria for small fibre neuropathy:
from symptoms to neuropathology

Grazia Devigili,¹ Valeria Tugnoli,² Paola Penza,³ Francesca Camozzi,³ Raffaella Lombardi,³ Giorgia Melli,³ Laura Broglio,³ Enrico Granieri¹ and Giuseppe Lauria⁴

¹Neurological Clinic, University of Ferrara, ²Neurophysiological Unit, S. Anna General Hospital, Ferrara, ³Neuromuscular Diseases Unit, National Neurological Institute 'Carlo Besta', Milan and ⁴Neurological Clinic, University of Brescia, Italy

Correspondence to: Dr Giuseppe Lauria, Neuromuscular Diseases Unit, National Neurological Institute 'Carlo Besta', via Celoria, 11, 20133 Milan, Italy
E-mail: glauria@istituto-besta.it

PRIMERA CONCLUSIÓN:

- En la AhTTR es frecuente el debut con afectación de FF en sus formas de inicio precoz >tardía.
 - RECORDAR:
 - NEUROPATÍA AUTONÓMICA.
 - NEUROPATÍA DE FIBRAS FINA (A+TermoAlgésica).
- PARA EL DIAGNÓSTICO PRECOZ y SEGUIMIENTO DE PORTADORES:
 - ¿QUÉ PRUEBAS NOS RECOMIENDAN HACER?

SEGUIMIENTO DE PORTADORES VAL30MET

Table 2 – ATTReuNET-recommended minimal monitoring of asymptomatic carriers and criteria for diagnostic confirmation of overt TTR-FAP.

History	Clinical questionnaire for sensory and autonomic dysfunction (gastrointestinal, erectile dysfunction, urinary dysfunction), SFN-SIQ
Body mass index (weight, height)	
Sensorimotor examination and testing	Temperature pain sensitivity in feet and legs
Neurophysiological testing	Neuropathy Impairment Score (NIS)
	Sweat test (Sudocan)
Autonomic examination and testing	Electromyography, nerve conduction studies (NCS)
	Quantitative sensory testing (QST) ^a
	Sympathetic skin response
	Heart rate response/variability ^b
Other	Microalbuminuria
Confirmation of diagnosis	Electrocardiography (ECG)
	Onset of symptoms and/or signs
	Changes in physiological tests compared with baseline
	Biopsy evidence of amyloid (confirmatory)

From Obici et al., 2016 [24].

ATTReuNET: European network for TTR-FAP or amyloidosis; TTR-FAP: transthyretin familial amyloid polyneuropathy (hereditary); SF-SIQ: short-form Symptom Inventory Questionnaire.

^a Required in Portugal.

^b Heart rate response to deep breathing.

Table 3 – Small-fiber neuropathy Symptom Inventory Questionnaire (SFN-SIQ): at least two symptoms are required to suggest the diagnosis of TTR-FAP in carriers of TTR mutation.

Sensory	0 or 1
Peripheral pain (burning)	
Paresthesia	
Allodynia	
Diminished temperature sensation	
Diminished pain sensation	
Autonomic	>2
Sicca syndrome (dry mouth/eyes)	
Accommodation disturbances	
Hyperhidrosis/hypohidrosis	
Micturition disturbances	
Impotence/diminished lubrication	
Gastroparesis	
Diarhea/constipation	
Facial flushing	
Orthostatic complaints	
Palpitations	
Total: 15	

From Lauría et al., 2012 [25].

TTR-FAP: transthyretin familial amyloid polyneuropathy (hereditary).

International meeting of the French society of neurology 2016

Familial amyloid polyneuropathy: When does it stop to be asymptomatic and need a treatment?

D. Adams^{a,b,d,*}, G. Beaudonnet^{b,e}, C. Adam^{b,c,d}, C. Lacroix^{b,c,d},
 M. Théaudin^{a,b}, C. Cauquil^{a,b}, C. Labeyrie^{a,b}

- PROTOCOLO DE ESTUDIO ANTE UNA SOSPECHA DE TTR-PNP.
- SEGUIMIENTO PORTADORES.

PRINCIPALES EXPLORACIONES NEUROFISIOLÓGICAS .

EXPLORACION NEUROFISIOLOGICA

- **EMG:** ENG /VCN $\Rightarrow$ exploramos fibra gruesa (M/S) (A α /A β).
- **Valoración de FIBRA FINA termo-algésica (A δ y c).**
 1. **QST (TEST SENSORIAL CUANTITATIVO).**
 2. LEPs (PE por laser).
 3. CHEPs (PE por calor).
 4. PREMPs. (PE por estímulo eléctrico doloroso) .
- **Valoración de FIBRA FINA autonómica (c).**
 1. **RESPUESTA SIMPATICO CUTANEA** (SN simpático): RSC.
 2. **INTERVALO R-R** (SN parasimpático): HrVRR.
- **MICRONEUROGRAFÍA (A δ y c) .**

1. ELECTROMIOGRAFÍA

- Ante la sospecha de neuropatía de fibra fina:
 - **ENG debe ser NORMAL** + soporte de test específicos.
- Ante la sospecha de inicio síntomas PNP en paciente portador :
 - **VALORAR CAMBIOS SIGNIFICATIVOS EN ENG BASAL**
(seguimiento evolutivo): caída de amplitud del SNAP/CMAP.
- Ante una sospecha de PNP HEREDITARIA POR TTR
esperaremos encontrar:
 - **POLINEUROPATIA SENSITIVA > MOTORA DE PREDOMINIO AXONAL** (amplitudes bajas) de inicio DISTAL EN EEII +STC.

2. VALORACIÓN NEUROFISIOLÓGICA DE FIBRA FINA: RESPUESTA SIMPÁTICO CUTÁNEA.

- Es el potencial registrado ante el **cambio de resistencia cutánea** provocado por la sudoración **tras un estímulo**.
 - ESTÍMULO SOMESTÉSICO, ACÚSTICO O MAGNÉTICO.
 - Información indirecta sobre la función sudomotora colinérgica simpática.

- **FISIOLOGÍA:**

- Reflejo polisináptico.
- Múltiples aferencias: estímulo emocional, acústico, eléctrico-sensitivo.
- SNC: hipotálamo, formación reticular, médula espinal (columna lateral) ipsilateral.
- Vía eferente: **fibras simpáticas sudomotoras pre y postganglionares (fibra amielínica C).**
- Activación glándulas sudoríparas. Está presente en pacientes con ausencia congénita de gl.sudoríparas.

Fig 1: A typical sympathetic skin response recording. Typical SSR recordings taken from the right palm (upper trace) and left palm (lower trace) (a) following electrical stimulus of left median nerve and (b) magnetic stimulation of cervical cord.

2. VALORACIÓN NEUROFISIOLÓGICA DE FIBRA FINA: RESPUESTA SIMPÁTICO CUTÁNEA.

ESPECIFICACIONES TÉCNICAS:

a) Estímulo eléctrico

- N.mediano, n.tibial, n.trigémino.
- Dur 0.1-0.2ms.
- Int 10-30mA (x3 umbral sens).
- No doloroso.
- Frecuencia irregular (>30sg)
 - Evitar habituación.

b) Estímulo acústico

c) Estímulo magnético, estímulo láser.

Registro

- Mano: palma-dorso mano.
- Pie: planta-dorso pie.
- Tiempo: 5-10s.
- Filtros: LFF 0,1-2Hz, HFF 100Hz.
- Sens: 0.2-2mV/div

Ventajas:

- * Fácil realización.
- * Método cribaje, screening.
- * **ALTA ESPECIFICIDAD.**

Desventajas:

- * Gran variabilidad.
- * Tendencia habituación.
- * No claros criterios definidos de anormalidad.
 - * ausencia
 - * ↓ amplitud >50% D-E
- * **BAJA SENSIBILIDAD.**

Mayor utilidad en la valorar la funcionalidad de fibras
fina (C) más que en patología autonómica.

2. VALORACIÓN NEUROFISIOLÓGICA DE FIBRA FINA: RESPUESTA SIMPÁTICO CUTÁNEA.

NORMAL

RSC PATOLOGICA PARA PIE

2. VALORACIÓN NEUROFISIOLÓGICA DE FIBRA FINA: RESPUESTA SIMPÁTICO CUTÁNEA.

years. As discussed above, the skin is controlled by conduction along unmyelinated fibers, and latency measurement is not useful. Initially regarded as an all-or-none phenomenon, now as techniques are refined, amplitude criteria are defined. According to Hoeldtke,³⁷ latency in hands is 1.6 ± 0.1 s, amplitude in hands is 1.3 ± 0.2 mV, latency in feet is 2.1 ± 0.1 s, and amplitude in feet is 0.8 ± 0.1 mV. According to Knezevic,⁴⁵ latency in hands is 1.5 ± 0.1 s, amplitude in hands is 0.5 ± 0.1 mV, latency in feet is 2.1 ± 0.2 s, and amplitude in feet 0.1 ± 0.04 mV. According to Drory,¹⁹ mean latency in hands is 1.5 s, mean amplitude in hands is 0.450 mV, mean latency in feet is 1. s, and mean amplitude in feet is 0.15 mV.

Clinical Comments. The main advantages of the SSR are that it is sensitive, reproducible, semiquantitative, simple, fast, and readily obtained on most electrophysiologic equipment. The main disorder

- >1.1 mV mano
- >0.7 mV pie

- >0.4 mV mano
- >0,15 mV pie

PATOLÓGICO

2. VALORACIÓN NEUROFISIOLÓGICA DE FIBRA FINA: INTERVALO R-R

- Registro de la variabilidad de la frecuencia cardiaca (R-R) en reposo.
 - Arritmia fisiológica sinusal (n.vago).
- Registro de la variabilidad de la frecuencia cardiaca (R-R) tras activaciones (HV, Valsalva).
- Programa automático (cálculo): media de 5 ciclos.

2. VALORACIÓN NEUROFISIOLÓGICA DE FIBRA FINA: INTERVALO R-R

a) R-R Intervalo. Condiciones basales.

- ❖ Registro durante 1 min.
- ❖ Promedio 5 min.
- ❖ Arritmia sinusal normal:
inspiración $\downarrow$ RR
expiración $\uparrow$ RR
- ❖ Variabilidad $>10\%$ RR

b) R-R Intervalo .

Respiración profunda 6 rpm.

- ❖ Registro durante 1 min a 6 rpm (5''I / 5''E).
- ❖ Inspiración $\downarrow$ RR - Espiración $\uparrow$ RR
- ❖ Diferencias entre FC min y FC max >10 lpm

2. VALORACIÓN NEUROFISIOLÓGICA DE FIBRA FINA: INTERVALO R-R

NORMAL

PATOLÓGICO

3. VALORACIÓN NEUROFISIOLÓGICA DE FIBRA FINA: QST (TEST CUANTITATIVO SENSORIAL).

- TEST PSICOFÍSICO (no invasivo) de evaluación de **UMBRALES SENSITIVOS** a estímulos sensoriales cuantificados en las diferentes vertientes sensitivas.
 - Exploración más generalizada para la detección de disfunción fibras finas.
 - Batería de pruebas.
- Permite evaluar de forma objetiva la SENSIBILIDAD y cuantificar la TERMOALGESIA.
- Permitirá evaluar tanto la **HIPOESTESIA COMO LA HIPERALGESIA.**
 - Distintas modalidades sensibilidad:
 - **térmica:** A δ , C (thermotest) –método de los límites-niveles-
 - vibratoria: A β (estímulo vibratorio-vibrometro-).
 - táctil: A β (monofilamentos).
- Estudio funcionalidad totalidad del circuito nervioso sensitivo : central y periférico.
- Pobre efecto localizador.
- Depende colaboración del paciente.
- **“Time consuming” ...**

3. VALORACIÓN NEUROFISIOLÓGICA DE FIBRA FINA: QST (TEST CUANTITATIVO SENSORIAL).

CASE IV

TSA II
Q SENSE

3. VALORACIÓN NEUROFISIOLÓGICA DE FIBRA FINA: QST (TEST CUANTITATIVO SENSORIAL).

BATERIA DE TEST en el QST :

1. UMBRALES SENSIBILIDAD TACTIL SUPERFICIAL:

- Monofilamento Semmes-Weinstein
- Filamentos de von Frey
- Distintos algoritmos
- PJ Dyck (WR Medical Electronics):
 - el test se inicia en la zona 2 y luego en la 8; según resultados obtenidos se utilizan distintos grosores de filamentos, y el test sigue distalmente, o bien proximalmente.
 - puntuación según resultados en cada punto anatómico. escala cuantitativa de deterioro neuropático.

2. UMBRALES SENSIBILIDAD VIBRATORIA:

- Estímulo vibratorio onda sinusoidal
 - Variación frecuencia vibración.
 - Variación intensidad (amplitud) vibración (125 Hz): 0-30g + frec.

Fibras A β

3. VALORACIÓN NEUROFISIOLÓGICA DE FIBRA FINA: QST (TEST CUANTITATIVO SENSORIAL).

3. UMBRALES TERMO-ALGESICOS (THERMOTEST):

- **Termodo:** aplicación sobre la piel.
 - **Principio de Peltier:** el paso de corriente eléctrica a través de dos semiconductores genera calor o frío dependiendo de la dirección de la corriente.
 - Temperatura adaptación: 32°C.
 - Rango temperaturas: 0°C – 50°C.
- **SE VALORAN UMBRALES AL:**
 - **Frío:** fibras A δ .
 - **Calor:** fibras C.
 - **Dolor por frío:** fibras A δ y C.
 - **Dolor por calor:** fibras C.
- **ÚTIL:** CUANTIFICA TANTO HIPOESTESIA COMO HIPERALGESIA.
- Variabilidad explorador/colaboración paciente.

3. VALORACIÓN NEUROFISIOLÓGICA DE FIBRA FINA: QST (TEST CUANTITATIVO SENSORIAL).

- UMBRALES: TÉCNICA:

- Estímulo dinámico. MÉTODO DE LOS LÍMITES.

- **Estímulo continuo** de intensidad creciente o descendente.
- El paciente señala cuando el estímulo es percibido.
- Umbral, tiempo de reacción incluido.

- Estímulo estático . MÉTODO DE LOS NIVELES.

- Varias **series de estímulos** estáticos de distinta magnitud .
- Cambio de la intensidad del estímulo.
- Estímulos nulos.
- Algoritmo 4,2,1: 15-20 estímulos
- Umbral: promedio.

- CUANTIFICACIÓN DEL DOLOR AL CALOR:

- Estudio perfil percepción dolor. Estímulo estático.
- Respuesta en base a una escala visual analógica (VAS):
- Cálculo: **Umbral HP:0.5** → inicio dolor. **Umbral HP:5.0** → nivel intermedio de dolor.

- La diferencia entre los umbrales HP:0.5 y HP:5.0 umbral muestra la tolerancia del sujeto a los estímulos dolorosos crecientes

MÉTODO DE LOS LIMITES: THERMOTEST

Test de Función SUDOMOTORA

- SUDOSCAN:
 - Cuantifica la conductancia del Cl⁻ (microsiemens, μ S) tras un estímulo eléctrico (placa acero) a 4v (desencadena respuesta electroquímica $\boxtimes$ secreción de Cl⁻ por las glándulas sudoríparas).
 - Sirve como biomarcador de la función de las gl.sudoríparas.
 - Localizar pacientes que tengan “riesgo de disfunción autonómica”.

II JORNADAS AMILOIDOSIS HEREDITARIA por TRANSTIRRETINA (A β TR)

Asociación Española
Enfermedad de Andrade
(Amiloidosis Hereditaria por Transtirretina)

Table 3. Diagnostic tests in small fiber neuropathy

Diagnostic test	Sensitivity	Specificity	Limitation	Sarcoidosis
SFNSL score <11 SFNSL score >48	100% 19%	31% 100%	Only validated in sarcoidosis	+
Skin wrinkling [59–62]	66–82%	67–75%	Training of staff necessary, no correlation with IENFD, poor NPV	No research available
QST (mainly TTT) [7,63–67]	57–93% TTT: 84% [63]	37–94% TTT:94% [63]	Requires conscious integration of patients, difficult to distinguish between faked and true loss of sensation, central and peripheral nervous system abnormalities can lead to same deficit	Sensitivity TTT 69%; abnormalities feet vs. hands (54 vs. 38%), 15% both [64]
LEP/CHEP/PREP [68–72,73**]	64–94%	69–87%	Abnormal test not only small fibers (entire nociceptive pathway), seasonal differences in heat conductivity in CHEP (less reliable outcome), limited availability, and needs well trained staff	No research available
Skin biopsy/IENFD [7,29,74,75]	33–90% Pure SFN: >90% [2,74]	58–95% Pure SFN: >90% [2,74]	Requires training, limited availability/not generally distributed (only a few centers), unclear in case of patchy disease, and time consuming	Sensitivity of 32.8% with age-dependent decrease (men < women) [29]
CCM/CNFD [76*,77*,78,79]	50–86%	52–84%	Not widely available, high acquisition costs, uncertainties about age-related decline, and racial differences in corneal nerve measurements	50% reduction in corneal nerve fibers in painful SFN compared with healthy controls [77*]
Autonomic function				
CAFT	Low			Abnormal in only 12.5% of SSFN patients [80]
QSART [66,81,82] Sudoscans [83,84]	59–80% 65–78%	– 80–92%	–	No research available on QSART or sudoscans. SSR abnormal in <10% of SSFN [64]
MIBG-scan	–	–	No standardized methodology or normative values, discontinuation of some drugs is needed	>50% involvement cardiac autonomic dysfunction [80]

CAFT, cardiovascular autonomic testing; CCM, corneal confocal microscopy; CHEP, contact heat-evoked potential; CNFD, corneal nerve-fiber density; IENFD, intraepidermal nerve-fiber density; LEP, laser-evoked potential; MIBG-scan, metaiodobenzylguanidine-scan; NPV, negative predictive value; PREP, pain-related evoked potential; QSART, quantitative sudomotor axon reflex test; QST, quantitative sensory testing; SFNSL, small fiber neuropathy screening list; SSFN, sarcoidosis small fiber neuropathy; SSR, sympathetic skin response; TTT, temperature threshold test.

Small fiber neuropathy: a disabling and underrecognized syndrome

Mareye Voortman^{a,b}, Daan Fritz^c, Oscar J.M. Vogels^d, Filip Eftimov^e, Diederik van de Beek^e, Matthijs C. Brouwer^e, and Marjolijn Drent^{a,e}

Curr Opin Pulm Med 2017, 23:447–457

ELSEVIER

Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com

ola
rade
Transthyretina

International meeting of the French society of neurology 2016

Familial amyloid polyneuropathy: When does it stop to be asymptomatic and need a treatment?

D. Adams^{a,b,d,*}, G. Beaudonnet^{b,e}, C. Adam^{b,c,d}, C. Lacroix^{b,c,d},
M. Théaudin^{a,b}, C. Cauquil^{a,b}, C. Labeyrie^{a,b}

^aService de neurologie, CHU Bicêtre, 78, rue du Général-Leclerc, 94275 Le Kremlin-Bicêtre cedex, France

^bCentre de référence national des neuropathies amyloïdes familiales et autres neuropathies périphériques rares (NNEFR), 94275 Le Kremlin-Bicêtre cedex, France

^cService d'anatomopathologie, CHU Bicêtre, 78, rue du Général-Leclerc, 94275 Le Kremlin-Bicêtre cedex, France

^dInsERM U1195, 78, rue du Général-Leclerc, 94275 Le Kremlin-Bicêtre cedex, France

^eUnité de neurophysiologie clinique et épileptologie, CHU Bicêtre, 78, rue du Général-Leclerc, 94275 Le Kremlin-Bicêtre cedex, France

Optimizing the management of transthyretin familial amyloid polyneuropathy in Europe: early diagnosis and effective care

David Adams, on behalf of the European Network for TTR-FAP (ATReuNET)*

Journal of the Peripheral Nervous System 21:5–9 (2016)

OPEN

Recommendations for presymptomatic genetic testing and management of individuals at risk for hereditary transthyretin amyloidosis

Laura Obici^a, Jan B. Kuks^b, Juan Buades^c, David Adams^d, Ole B. Suhr^e,
Teresa Coelho^f, Theodore Kyriakides^g, from the European Network for
TTR-FAP (ATReuNET)

NEUROPHYSIOLOGICAL TECHNIQUES TO DETECT EARLY SMALL-FIBER DYSFUNCTION IN TRANSTHYRETIN AMYLOID POLYNEUROPATHY

ISABEL CONCEIÇÃO, MD,^{1,2} JOÃO COSTA, PhD,² JOSÉ CASTRO, Tech,^{1,2} and MAMEDE DE CARVALHO, MD, PhD^{1,2}

¹Department of Neurosciences, Centro Hospitalar Lisboa Norte- Hospital de Santa Maria, Av Prof Egas Moniz, 1649-028, Lisbon, Portugal

²Translational and Clinical Physiology Unit, Instituto de Medicina Molecular, Faculty of Medicine, Lisbon, Portugal

Accepted 4 May 2013

REVIEW

“Red-flag” symptom clusters in transthyretin familial amyloid polyneuropathy

Isabel Conceição¹, Alejandra González-Duarte², Laura Obici³, Hartmut H.-J. Schmidt⁴,
Damien Simoneau⁵, Moh-Lim Ong⁶, and Leslie Amass⁶

¹CHLN – Hospital de Santa Maria, and Clinical and Translational Physiology Unit, Faculty of Medicine-IMM, Physiology Institute, Lisbon, Portugal; ²Department of Neurology, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, México City, México; ³Amyloidosis Research and Treatment Center, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy; ⁴Klinik für Transplantationsmedizin, Universitätsklinikum Münster, Münster, Germany; ⁵Medical Division, Pfizer International Operations, Paris, France; and ⁶Global Medical Affairs, Global Innovative Pharma, Pfizer Inc, New York, NY, USA

SEGUIMIENTO PORTADORES VAL30MET HSL

RESEARCH REPORT

Transthyretin familial amyloid polyneuropathy (TTR-FAP) in Mallorca: a comparison between late- and early-onset disease

Juan Buades-Reinés¹, Manuel Raya-Cruz¹, Cristina Gallego-Lezaún¹, Tomás Ripoll-Vera², Mercedes Usón-Martín³, Hernán Andreu-Serra⁴, and Eugenia Cisneros-Barroso⁵

¹Department of Internal Medicine, Son Llàtzer Hospital, Carretera de Manacor; ²Department of Cardiology, Son Llàtzer Hospital, Carretera de Manacor; ³Department of Neurology, Son Llàtzer Hospital, Carretera de Manacor; ⁴Department of Digestive Medicine, Son Llàtzer Hospital, Carretera de Manacor; and ⁵Clinical Research Coordinator A-TTR Group, Son Llàtzer Hospital, Carretera de Manacor, Palma, Spain

Table 1. Patient baseline demographics and disease characteristics.

Characteristics	Early AO (<50 years) (n = 53)	Late AO (≥50 years) (n = 42)	p-value
Mean ± SD age of onset, years	35 ± 9	64 ± 8	<0.001
Male/female	33/20	21/21	NS
Initial symptoms, n (%)			
Autonomic involvement	36 (68)	38 (90)	0.007
Sensory-motor	16 (30)	4 (9)	0.01
Cardiac involvement	1 (2)*	0 (0)	NS
Other organ involvement	23 ± 4	26 ± 4	0.02
Mean ± SD body mass index (kg/m ²)	23 ± 4	26 ± 4	0.02
Mean ± SD serum albumin (g/dl)	4 ± 0.5	3.7 ± 0.5	0.02

*, age of onset; NS, non-significant difference between early and late AO groups; SD, standard deviation.

Table 2. Clinical features during follow-up

Feature	Early AO (<50 years) n = 53 (%)	Late AO (≥50 years) n = 42 (%)	p-value
Neurologic involvement	45 (85)	36 (86)	NS
Autonomic dysfunction	39 (74)	35 (83)	NS
Digestive involvement	38 (72)	28 (67)	NS
Renal involvement	8 (15)	13 (31)	0.05
Cardiac involvement	–	16 (38)	NS
Syncope	12 (23)	5 (12)	0.01
Dyspnea	0 (0)	8 (19)	0.01
Chronic heart failure	2 (4)	10 (24)	0.004
Ophthalmologic involvement	9 (17)	13 (31)	NS
Dry eye	2 (4)	7 (17)	0.03
Vitreous opacity	7 (13)	12 (29)	0.05

AO, age of onset; NS, non-significant difference between early and late AO groups.

Neurophysiological markers in familial amyloid polyneuropathy patients: Early changes

Isabel M. Conceição ^{a,*}, José F. Castro ^a, Manuel Scotto ^b, Mamede de Carvalho ^a

^a Department of Neurology, Hospital de Santa Maria and Neuromuscular Unit, Institute of Molecular Medicine, Faculty of Medicine, Av Prof Egas Moniz, 1649-028 Lisbon, Portugal

^b Department of Mathematics, University of Aveiro, Aveiro, Portugal

Accepted 10 January 2008
Available online 4 March 2008

Abstract

Objective: Familial amyloid polyneuropathy-type I (FAP-I) is a hereditary, axonal, sensory-motor and autonomic polyneuropathy, with early involvement of small fibres. Liver transplantation is the only effective therapy in FAP, but should be performed early in the course of the disease. Reliable quantitative methods that could allow the determination of early changes in the peripheral nerve function are essential. Our aim was to find sensitive neurophysiological markers in FAP-I.

Methods: Eighty-one FAP-I patients were included in this study. They were divided into two groups (G1, asymptomatic FAP-I mutation carriers; G2, early symptomatic). Seventy-six healthy controls formed a control group (G3). Nerve conduction studies, needle electromyography with motor unit potential analysis of the extensor digitorum brevis, RR interval and sympathetic skin response (SSR) were analyzed.

Results: The amplitudes of the motor response of the peroneus nerve and of the plantar SSR were significantly lower in G1 compared to G3. No other differences were found between those two groups. With a cut-off point of 0.2 mV for plantar SSR, its sensitivity and specificity are 0.53 and 0.95, respectively. The positive predictive value and the negative predictive value are 0.82.

Conclusions: SSR response at foot is a useful measurement to detect early dysfunction of peripheral nerve fibres in FAP-I. Its abnormality should be considered a warning sign and lead to a careful clinical assessment.

Significance: SSR is a useful neurophysiological marker in FAP-I.

© 2008 International Federation of Clinical Neurophysiology.

Published by Elsevier Ireland Ltd. All rights reserved.

Table 3	PORTADORES	SINTOMATICOS	SANOS			
Autonomic evaluation						
Test	G1 (n = 38)	G2 (n = 17)	G3 (n = 38)	G1 × G3	G1 × G2	G2 × G3
SSR						
Hand – Ampl (mV)	2.5 ± 1.6	0.9 ± 0.8	3.1 ± 2.1	NS	<0.01	<0.01
Hand – Latency (s)	1.4 ± 0.2	1.5 ± 0.1	1.4 ± 0.1	NS	NS	<0.01
Foot – Ampl (mV)	0.8 ± 0.6	0.3 ± 0.4	1.5 ± 1.1	<0.01	<0.01	<0.01
Foot – Latency (s)	2.2 ± 0.3	2.3 ± 0.3	2.1 ± 0.2	NS	NS	NS
	G1 (n = 31)	G2 (n = 12)	G3 (n = 36)	G1 × G3	G1 × G2	G2 × G3
RRIV						
Rest (%)	20.8 ± 9.8	10.7 ± 7.1	17.5 ± 7.6	NS	<0.01	NS
Deep breathing (%)	33.3 ± 15.5	21.6 ± 17.9	29.9 ± 14.0	NS	<0.01	NS

G, group; SSR, skin sympathetic response; RRIV, RR interval.
NS, non significant. Significant statistical comparisons are in bold.

- 1.ENG
- 2.EMG
- 3.RR
- 4.RSC

VALORES MENORES DE 0.2 mV de la RSC en pie deben ser considerados un SIGNO DE ALARMA.

NEUROPHYSIOLOGICAL TECHNIQUES TO DETECT EARLY SMALL-FIBER DYSFUNCTION IN TRANSTHYRETIN AMYLOID POLYNEUROPATHY

ISABEL CONCEIÇÃO, MD,^{1,2} JOÃO COSTA, PhD,² JOSÉ CASTRO, Tech,^{1,2} and MAMEDE DE CARVALHO, MD, PhD^{1,2}

¹Department of Neurosciences, Centro Hospitalar Lisboa Norte- Hospital de Santa Maria, Av Prof Egas Moniz, 1649-028, Lisbon, Portugal

²Translational and Clinical Physiology Unit, Instituto de Medicina Molecular, Faculty of Medicine, Lisbon, Portugal

Accepted 4 May 2013

Table 2. Diagnostic accuracy of the neurophysiological tests.*

Controls vs.	Plantar SSR Amp	LEP N2 Lat
TTR-FAP mutation carriers		
Specificity (%)	94 (80–98)	97 (85–99)
Sensitivity (%)	22 (11–39)	25 (13–41)
Diagnostic OR	4.3 (0.8–22.8)	10.7 (1.2–91.1)
AUC-ROC	0.65 (0.52 to 0.79)	0.65 (0.52 to 0.79)
Early symptomatic patients		
Specificity (%)	94 (80–98)	97 (85–99)
Sensitivity (%)	32 (19–50)	22 (9–45)
Diagnostic OR	7.4 (1.5–37.2)	9.1 (0.9–89.3)
AUC-ROC	0.75 (0.63 to 0.86)	0.76 (0.63 to 0.89)

AUC-ROC, area under the receiver operating characteristic curve; LEP N2 Lat, laser evoked potential latency; OR, odds ratio; SSR Amp, sympathetic skin response amplitude.

*Values in parentheses represent 95% confidence intervals. The cutoff values used for diagnostic accuracy analysis corresponded to the 5th percentile of the data distribution: 0.50 mV for plantar SSR Amp and 279 ms for LEP N2 Lat.

ABSTRACT: *Introduction:* Transthyretin familial amyloid polyneuropathy (TTR-FAP) is characterized by early selective involvement of small nerve fibers. Initial clinical diagnosis is complicated by psychosocial factors. We evaluated diagnostic accuracy of sural sensory nerve action potentials, plantar sympathetic skin response (SSR), and cortical laser-evoked potentials (LEP) to dorsal foot stimulation in the early diagnosis of TTR-FAP. *Methods:* Sixty-three subjects with TTR-FAP (Val30-Met) mutation were split into 2 groups (asymptomatic carriers and early-symptomatic patients) and compared with 33 healthy controls. *Results:* The diagnostic accuracy of plantar SSR amplitude and LEP N2 latency was similar; all had very high specificity (94 to 97%) but low sensitivity (22 to 32%) in distinguishing controls from carriers and early-symptomatic patients. No control had abnormal results on both tests. *Conclusions:* Plantar SSR and LEPs have similar diagnostic performance in detecting small-fiber dysfunction in early TTR-FAP; we propose that both tests should be used to investigate this population.

Muscle Nerve 49: 181–186, 2014

- 1.RSC
- 2.LEP
- 3.ENG

RESTO DE CONCLUSIONES:

- La NEUROPATÍA DE FIBRA FINA SE DIAGNOSTICA POR LA **CLINICA + TEST DIAGNOSTICO** que valore la FIBRA FINA NERVIOSA.
 - ALODINIA y/o HIPERESTESIA y/o ALTERACION PERCEPCION TERMICA.
 - QST patológico.
 - Biopsia cutánea.
- Para el seguimiento de portadores y diagnóstico precoz de la NEUROPATIA DE FIBRA FINA y PNP nos recomiendan realizar una **batería de test** neurofisiológicos que incluyen: **EMG, RSC, RR, QST.**
- HAY QUE “**PERSEGUIR**” SÍNTOMAS AUTONÓMICOS Y/O DE FIBRA FINA TERMOALGÉSICA que a pesar de ser en ocasiones INESPECIFICOS Y VARIABLES pueden ser estudiados por diferentes TEST QUE EXPLORAN DIFERENTES VERTIENTES DEL SNA y FF y QUE entre ellos SE COMPLEMENTAN (**buscar > 2 test alterados** : SNA o S.Termo-Algésica).
- ESTUDIO EMG/ENG ES NORMAL en la NEUROPATIA DE FIBRA FINA : un EMG normal no implica que no haya afectación de fibras nerviosas.

II JORNADAS
AMILOIDOSIS HEREDITARIA
por TRANSTIRRETINA (A α TTR)

Asociación Española
Enfermedad de Andrade
(Amiloidosis Hereditaria por Transtirretina)

GRACIAS