MANEJO DE LA ANTIAGREGACIÓN Y LA ANTICOAGULACIÓN EN LOS PROCEDIMIENTOS QUIRÚRGICOS

Juan V. Llau, MD, PhD

Dept. Anestesia y Reanimación – Hospital Clínico de Valencia

Profesor Asociado de Fisiología - Universidad de Valencia

juanvllau@gmail.com

Servicio de Anestesiología,
Reanimación y
Terapéutica del Dolor

Hospital Clínico Universitario de Valencia

Conflicto de intereses

Conflicto de intereses

Agradecimiento especial a todos aquellos que forman parte de mi excelente equipo de colaboradores y a todos aquéllos que confían en mi para compartir con vosotros nuestro trabajo

Inicio

ANTICOAGULANTS	ORAL ANTICOAGULANTS	VKA: Warfarin, acenocoumarol
		NOAC: Apixaban, Rivaroxaban, Dabigatran
	PARENTERAL ANTICOAGULANTS	DIRECT: Desirudin
		INDIRECT: UFH, LMWH, Fondaparinux
ANTIPLATELET AGENTS	REVERSIBLE ACTION	COX-INHIBITOR: NSAID
		COX-INHIBITOR: Aspirin
	IRREVERSIBLE ACTION	ADP-r ANTAGONISTS: Thienopyridins

Punto de partida: estratificación

Review Article

The perioperative management of new direct oral anticoagulants: a question without answers

Raquel Ferrandis¹; Jordi Castillo²; José de Andrés³; Carmen Gomar⁴; Aurelio Gómez-Luque⁵; Francisco Hidalgo⁶; Juan V. Llau⁷;
Pilar Sierra⁸; Luis M. Torres⁹

Thromb Haemost 2013; 110: 515–522

**RIESGO
TROMBÓTICO**

**RIESGO
HEMORRÁGICO**

**RIESGO
"RENAL"**

Punto de partida: estratificación

RIESGO TROMBÓTICO

Review Article

The perioperative management of new direct oral anticoagulants: a question without answers

Raquel Ferrandis¹; Jordi Castillo²; José de Andrés³; Carmen Gomar⁴; Aurelio Gómez-Luque⁵; Francisco Hidalgo⁶; Juan V. Llau⁷; Pilar Sierra⁸; Luis M. Torres⁹

Thromb Haemost 2013; 110: 515–522

RIESGO

Low	Moderate	High
Atrial fibrillation		
CHA ₂ DS ₂ -VASc 0–1 points No other risk factor	CHA ₂ DS ₂ -VASc 2–4 points	CHA ₂ DS ₂ -VASc >4 points Stroke within 3 months Rheumatic valvulopathy

Bajo

Moderado

Alto

Riesgo trombótico (tratamiento con antiagregantes)

Más de 6 meses tras IAM, cirugía coronaria, coronariografía percutánea, *stent* metálico o ACV (más de 12 meses en caso de complicaciones)

Entre 6 y 24 semanas tras IAM, cirugía coronaria, coronariografía percutánea, *stent* metálico o ACV (6-12 meses en caso de complicaciones o DM o baja FE). Más de 12 meses tras *stent* farmacológico

Menos de 6 semanas tras IAM, cirugía coronaria, coronariografía percutánea, *stent* metálico o ACV (menos de 6 meses si complicaciones). Menos de 12 meses tras *stent* farmacológico

Punto de partida: estratificación

Review Article

The perioperative management of new direct oral anticoagulants:
a question without answers

Raquel Ferrandis¹; Jordi Castillo²; José de Andrés³; Carmen Gomar⁴; Aurelio Gómez-Luque⁵; Francisco Hidalgo⁶; Juan V. Llau⁷;
Pilar Sierra⁸; Luis M. Torres⁹

Thromb Haemost 2013; 110: 515–522

**RIESGO
HEMORRÁGICO**

**RIESGO
TROMBÓTICO**

PERIOPERATORIO

Tipo de cirugía
Tipo de anestesia

Tiempo de suspensión
Bridging

ESPONTANEO

HAS-BLED

Edad

Creatinina

Otros por definir

**RIESGO
"RENAL"**

Punto de partida: estratificación

Review Article

The perioperative management of new direct oral anticoagulants: a question without answers

Raquel Ferrandis¹; Jordi Castillo²; José de Andrés³; Carmen Gomar⁴; Aurelio Gómez-Luque⁵; Francisco Hidalgo⁶; Juan V. Llaú⁷; Pilar Sierra⁸; Luis M. Torres⁹

Thromb Haemost 2013; 110: 515–522

**RIESGO
"RENAL"**

**RIESGO
TROMBÓTICO**

**RIESGO
HEMORRÁGICO**

ACLARAMIENTO DE CREATININA

Fórmula de Crockcoft-Gault

$$\text{CrCl} = [(140 - \text{edad}) \times \text{peso}] / (72 \times \text{Cr})$$

(en mujeres x 0.85)

SIN RIESGO	RIESGO BAJO	RIESGO MODERADO	RIESGO ALTO
CrCl > 80	CrCl 50-80	CrCl 30-50	CrCl < 30

ACOD

Available online at
SciVerse ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com/en

ACOD

GUIDELINES

Surgery and
long-term
Thrombin
of the Working
and the
haemostasis

Management of major bleeding complications and emergency surgery in patients on long-term treatment with direct oral anticoagulants, thrombin or factor-Xa inhibitors: Proposals of the Working Group on Perioperative Haemostasis (GIHP) – March 2013

Chirurgías et actes invasifs chez les patients traités au long cours par un anticoagulant oral anti-IIa ou anti-Xa direct. Recommandations du Groupe d'intérêt en hémostase périopératoire (GIHP) et du Groupe d'études sur l'hémostase et les...

s // Cons
EIO PE
AGULA
REUNIÃO DE C
CRISTIANA FONSECA¹, JOANA ALVES²,

bjh guideline

Guideline on the management of bleeding in patients on antithrombotic agents

Mike McBride,^{1,2} Janet J. Van Viers,³ Geoffrey B. Tait,⁴ Andrew D. Woodford,⁵ Mike Laffan⁶ and on behalf of the British Committee for Standards in Haematology

DO DA 3ª
LOGIA

Managing New Oral Anticoagulants in the Perioperative and Intensive Care Unit Setting

Jerrold H. Levy, M.D., FAHA, F.C.C.M.,* David Faraoni, M.D.,† Jenna L. Spring, M.S.,‡ James D. Douketis, M.D.,§ Charles M. Samama, M.D., Ph.D., F.C.C.P.¶

direct oral anticoagulants in the perioperative

Acute management of bleeding in patients on novel oral anticoagulants

Deborah M. Siegal¹ and Mark A. Crowther^{1,2*}

Division of Hematology and Thrombocytosis, Department of Medicine, McMaster University, Hamilton, Ont., Canada and ²Subacute Medicine, St Joseph's Healthcare and Hamilton Health Sciences, Hamilton

Received 18 June 2012; revised 19 September 2012; accepted 11 October 2012

Revista Española de Anestesiología

ARTÍCULO ESPECIAL

Manejo de los anticoagulantes orales de acción directa en el periodo perioperatorio y técnicas invasivas

J.V. Llau^{a,*}, R. Ferrandis^b, J. Castillo^c, J. de Andrés^d, C. Gomar^e, A. Gómez-Luque^f, F. Hidalgo^g y L.M. Torres^h, en representación de los participantes en el Foro de Consenso de la ESRA-España sobre «Fármacos que alteran la hemostasia»^g

Guidance on the emergent reversal of oral thrombin and factor Xa inhibitors

Scott Kaatz,^{1*} Peter A. Kouides,² David A. Garcia,³ Aley C. Spyropoulos,⁴ Mark Crowther,⁵ Jim D. Douketis,⁶ Anthony K. C. Chan,⁷ Andra James,⁸ Stephan Möl,⁹ Thomas L. Ornt,¹⁰ Elizabeth M. Van Cott,¹¹ and Jack Ansell¹¹

**Perioperative Management
of Patients Receiving New
Oral Anticoagulants**
An International Survey

David Faraoni, MD^{1,2}, Charles Marc Samama, MD, PhD³,
Marco Ranucci, MD, FRC⁴, Wulf Dietrich, MD, PhD⁵,
Jerrold H. Levy, MD, FAHA, FCCM⁶

ACOD-perioperatorio

Clin Lab Med 34 (2014) 637–654

Table 3

Articles on recommendations for the perioperative management of patients treated with NOACs

Author, Year	Endorsed	Comments
Rosencher et al, ²⁹ 2007		Regional anesthesia
Llau & Ferrandis, ²⁸ 2009		Regional anesthesia
Gogarten et al, ²⁷ 2010	ESA	Regional anesthesia
Horlocker et al, ²⁶ 2010	ASRA	Regional anesthesia
Levy et al, ²⁵ 2010		Review article
Sie et al, ¹² 2011	GIHP	Perioperative recommendations
Vandermeulen et al, ²⁴ 2011	BARA	Regional anesthesia
Spyropoulos et al, ⁹ 2012		Review article with recommendations
Levy et al, ¹⁰ 2013		Review article with recommendations
Kozek-Langenecker et al, ¹³ 2013	ESA	ESA guidelines
Spahn et al, ¹⁴ 2013		Updated European trauma guidelines
Steiner et al, ¹⁵ 2013	DKG	Emergency management
Fawole et al, ¹¹ 2013		Review article with recommendations
Ferrandis et al, ¹⁶ 2013	WGSFAA	Review article with recommendations
Pernod et al, ¹⁷ 2013	GIHP	Bleeding complications and emergency surgery

PERIPROCEDURAL MANAGEMENT OF DIRECT ORAL ANTICOAGULANTS:
CURRENT STATUS.

R. Ferrandis, J.V. Llau, J. De Andrés, C. Gomar, A. Gómez-Luque, F. Hidalgo,
P. Sierra, L.M. Torres, on behalf of Spanish Forum on Anticoagulants and
Anesthesia.

ACOD-perioperatorio

Pendiente de publicación

Fármaco CrCl	RIVAROXABAN APIXABAN		DABIGATRAN	
	>50	30-50	>50	30-50
Riesgo hemorrágico bajo	1 día	2 días	2 días	3 días
Riesgo hemorrágico moderado-alto	2 días	3 días	3 días	4 días
Riesgo trombótico alto	VALORAR TERAPIA PUENTE			

Abstracts and Highlight Papers of the 33rd Annual European
Society of Regional Anaesthesia & Pain Therapy (ESRA)
Congress 2014: Invited Speaker Highlight Papers

ESRA1-0676

Panel Discussion: Regional anaesthesia in the anticoagulated
patient – when allowed, when contraindicated

OLD AND NEW ANTICOAGULANTS: WHAT ARE THE
GUIDELINES SAYING?

Ferrandis Comes R.¹, Llau Pitarch J.V.¹ *Anaesthesiology and Critical Care,*
Hospital Clinic Universitari, Valencia, Spain.

ACOD-sangrado

CLINICAL RESEARCH

Management of major bleeding complications and emergency surgery in patients on long-term

tr
OI
G
2)
Pr
ur
di
mi

Haemorrhage and DABIGATRAN (PRADAXA®) or RIVAROXABAN (XARELTO®)

Bleeding into a critical organ (intracranial, acute subdural, intraocular...)

1) FEIBA® 30-50 UI / kg* or
2) PCC 50 UI / kg*

Haemorrhage and DABIGATRAN (PRADAXA®) or RIVAROXABAN (XARELTO®)

Serious bleeding according to the French Health Authority (2008) (excluding previous cases)

Bleeding into a critical organ (intracranial, acute subdural, intraocular...)

1) FEIBA® 30-50 UI / kg* or
2) PCC 50 UI / kg*

- If ratio aPTT ≤ 1.2 and ratio PT ≤ 1.2: no reversal
- Prefer haemostatic procedure if feasible
- If no haemostatic procedure is appropriate and if ratio aPTT > 1.2 (isolated) or ratio PT > 1.2
- ▶ Discuss reversal** (not always necessary) and obtain specific dosage

Serious bleeding according to the French Health Authority (2008) (excluding previous cases)

* Depending on availability. No data available on the thrombotic risk of high doses of PCC or FEIBA in these patients
** [] means concentration
*** PCC=25-50 UI/kg or FEIBA=30-50 UI/Kg
rFVIIa is not considered first-line

Antiagregantes plaquetarios

Manejo de los AAP

EJA

Eur J Anaesthesiol 2013; 30:1–112

GUIDELINES

Management of severe perioperative bleeding

Guidelines from the European Society of Anaesthesiology

Sibylle A. Kozek-Langenecker, Arash Afshari, Pierre Albaladejo, Cesar Aldecoa Alvarez Santullano, Edoardo De Robertis, Daniela C. Filipescu, Dietmar Fries, Klaus Görlinger, Thorsten Haas, Georgina Imberger, Matthias Jacob, Marcus Lancé, Juan Llau, Sue Mallett, Jens Meier, Niels Rahe-Meyer, Charles Marc Samama, Andrew Smith, Cristina Solomon, Philippe Van der Linden, Anne Juul Wikkelsø, Patrick Wouters and Piet Wyffels

ASPIRINA: Mantener en la mayoría de pacientes
Si se considera suspenderla: 5 días

CLOPIDOGREL: Suspender 5 días.
Valorar sustituir por AAS

PRASUGREL: Suspender 7 días

TODOS: Reinicio lo antes posible

Agent	Recommended Interval between Last Dose and Procedure
Antiplatelet agents	
Aspirin	
Aspirin and dipyridamole (Aggrenox, Boehringer Ingelheim)	2 days
Cilostazol (Pletal, Otsuka Pharmaceutical)	
Thienopyridine agents (clopidogrel [Plavix, Sanofi Aventis], ticlopidine [Ticlid, Roche], prasugrel [Effient, Eli Lilly], and ticagrelor [Brilinta, AstraZeneca])	5 days (clopidogrel and ticagrelor), 7 days (prasugrel), or 10–14 days (ticlopidine)

Manejo de los AAP

Manejo de los AAP

CIRUGÍA URGENTE

CIRUGÍA URGENTE SIN HEMORRAGIA ASOCIADA

- 1.- No administrar plaquetas de forma profiláctica
- 2.- Inicio de cirugía
- 3.- Si hemorragia durante la cirugía: plaquetas y manejo de la misma con CH + prohemostáticos

CIRUGÍA URGENTE CON HEMORRAGIA ASOCIADA

- 1.- Administrar plaquetas (1 pool de inicio)
- 2.- Inicio de cirugía
- 3.- Si hemorragia no controlable: plaquetas y manejo de la misma con CH + prohemostáticos

AVK

Manejo de los AVK

“Terapia Puente” VS “No Terapia Puente”

CHEST

Supplement

ANTITHROMBOTIC THERAPY AND PREVENTION OF THROMBOSIS, 9TH ED: ACCP GUIDELINES

Perioperative Management of Antithrombotic Therapy

Antithrombotic Therapy and Prevention of Thrombosis,
9th ed: American College of Chest Physicians
Evidence-Based Clinical Practice Guidelines

James D. Douketis, MD, FCCP; Alex C. Spyropoulos, MD, FCCP; Frederick A. Spencer, MD;
Michael Mayr, MD; Amir K. Jaffer, MD, FHM; Mark H. Eckman, MD;
Andrea S. Dunn, MD; and Regina Kunz, MD, MSc (Epi)

Results: In patients requiring vitamin K antagonist (VKA) therapy, we recommend **stopping VKAs 5 days before surgery instead of 10 days before surgery**. In patients with a mechanical heart valve, atrial fibrillation, or a recent stroke, we suggest **bridging anticoagulation instead of no bridging** (Grade 2C); in patients at low risk, we suggest **no bridging**. In patients who require a **dental procedure**, we suggest **bridging anticoagulation**.

• **Bridging dose regimens.** Three dose regimens have been studied:

1. A **high-dose (therapeutic-dose)** heparin bridging regimen involves administering an anticoagulant dose that is similar to that used for the treatment of acute VTE or an acute coronary syndrome (eg, enoxaparin 1 mg/kg bid or 1.5 mg/kg daily, dalteparin 100 International Units/kg bid or 200 International Units/kg daily, tinzaparin 175 International Units/kg daily, IV UFH to attain an activated partial thromboplastin time [aPTT] 1.5 to 2 times the control aPTT).
2. A **low-dose (prophylactic-dose)** heparin regimen involves administering a dose that is used, typically, to prevent postoperative VTE (eg, enoxaparin 30 mg/kg bid or 40 mg/kg daily, dalteparin 5,000 International Units/kg daily, UFH 5,000-7,500 International Units/kg bid).
3. An **intermediate-dose regimen** has recently been studied for bridging and is intermediate in anticoagulant intensity between high- and low-dose regimens (eg, enoxaparin 40 mg/kg bid).

Periprocedural Management of Patients on Anticoagulants

Lance A. Williams II, MD^{1,2}, James M. Hunter Jr, MD²,
Marika B. Mangano, MD¹, Thomas B. Victor, MD, PhD²

Clin Lab Med 34 (2014) 595–611

Manejo de los AVK

Conclusiones

El manejo y gestión de los AAP y los AC es un gran reto para el equipo anestésico-quirúrgico

AAS debe mantenerse si es posible hasta el día antes de la cirugía a dosis bajas (100 mg) y el clopidogrel debe sustituirse por AAS

En pacientes de alto riesgo trombótico, la doble antiagregación debe manejarse bajo un criterio multidisciplinar

El manejo óptimo de los ACOD en el periodo perioperatorio un protocolo para alcanzar el mejor balance de los riesgos trombótico y hemorrágico en cada paciente

El manejo de los AVK sigue siendo controvertido, aunque la tendencia actual parece ser terapia puente con dosis bajas de HBPM

MANEJO DE LA ANTIAGREGACIÓN Y LA ANTICOAGULACIÓN EN PROCEDIMIENTOS QUIRÚRGICOS

MUCHAS GRACIAS POR VUESTRA ATENCIÓN

Juan V. Llau, MD, PhD

Dept. Anestesia y Reanimación – Hospital Clínico de Valencia

Profesor Asociado de Fisiología - Universidad de Valencia

juanvllau@gmail.com